

WEED MANAGEMENT 2007

Prepared by Hilary A. Sandler

NEW HERBICIDES. I anticipate that at least one new herbicide should become registered after the printing of the 2007 Chart Book. Check our web site and upcoming newsletters for further information about these new products.

WEED LIFE CYCLES. *ANNUAL PLANTS* complete their life cycle in one year and must reproduce by seed. *PERENNIAL PLANTS* can live for many years and may reproduce by seed, runners, rhizomes, etc. Most of the weeds in cranberry production are perennials. With the exception of dodder, annual weeds are much less common and easier to control than perennials. However, infestations of annual weeds should not be taken lightly, especially on new plantings. Annual plants are designated with an (A). Unless otherwise noted, all other weeds are perennials.

WEED MAPPING. Weed maps can help you organize the management of your weed problems, especially with perennial weeds. Weed maps should be done every few years, depending on weed pressure and management objectives. Several steps are involved: 1) Correctly identify the weeds, 2) Document the location of the weeds (by drawings or photographs), and 3) Designate a priority number to the weed. Weeds are grouped into priorities (1=no tolerance; 2=serious concern; 3=less concern; and 4=lowest concern) based on their likelihood to cause yield loss, ability to spread, and the difficulty of control. Deal with the most yield-threatening weeds first. Recommended priority groupings are noted in the right-hand margin for each weed. Growers may change and adapt weeds into priority groups to best fit their own management program.

CLIPPER APPLICATORS. Roundup products are the only products currently labeled for use in clipper application (Section 2ee-'Cut Stump'). No other glyphosate products may be used in clipper applicators. Using the correct technique is critical for maximizing the performance of this herbicide application. Roundup must be applied to the stem as it is cut! Good stem coverage and adequate flow without dripping on the vines is essential. Concentrated solutions (50-100% Roundup) work best. Notations are made within each weed management description below if field studies have demonstrated clippers to be effective in giving partial or good weed control.

TIMING OF PREEMERGENCE HERBICIDES. Spring applications are typically done from March through mid-April. Fall applications are typically done 1-2 weeks after harvest but at least 2-3 weeks prior to the winter flood. Applications for dodder control are put out as close as possible to the time of seedling emergence (usually mid-April through mid-May).

DODDER (A) *Cuscuta gronovii*

PRIORITY 1

Prevention. The best management strategy for dodder control is prevention of infestation. This is best accomplished through good sanitation. Dodder seed is easily moved in harvest water and on equipment. When possible, dodder-infested beds should be harvested after clean beds. Floats used to corral berries should be inspected for the presence of dodder seed before they are placed in beds that are dodder-free. This is especially important for custom harvest operations. A good trash flow after harvest is also helpful in removing seed capsules from the bed, but is not a replacement for prevention since subsequent infestation occurs.

Dodder may also be spread vegetatively: pieces of stem can be moved on equipment, such as granular spreaders, and become attached to healthy plants. Care should be taken when moving any piece of equipment from a bed infested with dodder to one that is not.

Dodder is an obligate parasite and must have a host plant to survive. Early in the growing season, cranberry stems are somewhat resistant to dodder infection, so control of soft fleshy weeds that can act as hosts, such as goldenrod, loosestrife and asters, is important for early season control. Dodder emergence may continue for 2 to 3 months, which is far longer than the residual activity of most preemergence herbicides. Therefore, control of other hosts early in the season is especially important. Where dodder infestations are just beginning, careful scouting and hand removal of seedlings prior to infection is a good practice. Weeds that are infected with dodder should be completely removed from the bed, since dodder stems will re-grow from the portion that is embedded in the weed.

Scouting. Scout in areas where infestation has occurred in the past and in bare areas in beds infested previously with dodder. Often, dodder seed will accumulate in the areas where berries are removed from the beds at harvest.

22 Weeds

Begin scouting for dodder in early-mid April (unless your bog history or unusual weather conditions indicates otherwise). Mild winter temperatures may promote earlier emergence of dodder. Recent research indicates that dodder populations in MA emerge slowly at first, but then quickly peak and subside. In these studies, 50% and 90% of the seedlings emerged from about 30 and 46 days after first emergence, respectively. In addition, seedlings emerge later and later as the seeds accumulate years in the soil, creating overlapping generations. The most vigorous population is the one most recently seeded (from last year). It is not known how variable dodder populations are from bog to bog but it may reasonably be so. Growers should consider previous successful experiences, along with scouting, and plan applications of preemergence herbicides accordingly.

CHEMICAL RECOMMENDATIONS FOR DODDER CONTROL

Casoron 4G	30-60 lb/A	Apply within 10-14 days of first seedling emergence. Follow with 0.2" water to incorporate herbicide. Split applications can be used (up to a total of 100 lb/A in a 12-month period). Allow at least 3 weeks between applications. See Notes on Casoron.
-------------------	------------	---

Other Herbicide Considerations. In order to maximize effectiveness of Casoron, monitor the soil temperature with a standard soil temperature probe. The soil should be at least 60°F to allow Casoron to volatilize and work effectively. If frequent frost events and/or spring rains occur after application and prior to the volatilization of the herbicide, the water may move some herbicide below the dodder seed zone and reduce effectiveness of the herbicide. To avoid injury, do not apply as vines approach bud break.

Spring floods. Recent research and grower experience has shown that short (24-48 hr) floods in early to mid-May may be effective for reducing dodder infestations. Timing and temperature of the flood may affect efficacy. Research is on-going to refine recommendations. Floods should cover vine tips adequately. No adverse vine or yield impact has been reported. Dodder floods may coincide with floods used to control black-headed fireworm.

Postemergence Control. For light to moderate dodder infestations, raking is not recommended. Yield and quality are reduced and dodder growth is unaffected. For heavy infestations, raking can prevent the onset of upright dieback caused by infection. It can also prevent leaf-drop due to shading of the cranberry by the dodder canopy. Raking prior to seed set is not as effective in reducing dodder seed production as raking after seed set. Raking should be done before the seed capsules begin to dry. No benefit is provided from raking earlier as the dodder stems re-grow from the portions embedded within the cranberry. No benefit is gained from raking more than once.

Research is on-going to refine recommendations for postemergence (biological) control; check with the weed specialist for information. A good trash flow after harvest is also helpful in removing seed pods from the bog.

BRISTLY & PRICKLY DEWBERRY *Rubus hispida*, *R. flagellaris* (Running brambles) PRIORITY 1

Dewberries spread rapidly on bogs by rooting at the tips of canes. Both types will kill vines if allowed to spread. The most effective way to manage dewberries is to eliminate them as they invade the bog. Remove young plants by pulling or digging out by the roots. Chemical control of established plants is difficult because the weed grows close to vine level. Sparse or moderately colonized spots can be hand-wiped with glyphosate products. Clipping stems with Roundup-dispensing applicators may offer partial control.

Late water floods reduce numbers of dewberry crowns and offspring plants in that year. Summer refloods (May 10-12 through July 15-20) may be used in desperate circumstances. This will eliminate the crop for that year! It is particularly helpful to hand-pull or hand-wipe remaining dewberry plants after the flood has been withdrawn. Fall floods may suppress populations of *R. hispida*; results have varied from no impact to reduced crown density. Start the flood as early as possible (mid-September) and hold for 4 weeks for best dewberry management. Consider starting a fall flood in early-mid September, if possible. It is important to remove the flood by early November to allow the vines time to become dormant prior to winter.

Knife-raking or pruning in the fall may help uproot offspring plants. Significant dewberry patches should be scraped and replanted with new vines.

DEWBERRIES - CHEMICAL RECOMMENDATIONS

Roundup WeatherMAX	Mix 1 part glyphosate with 4-9 parts water (10-20% solutions). No additional additives, buffers, or surfactants are needed. However, the addition of ammonium sulfate may sometimes improve performance.
+ A marker dye (i.e., Blazon Blue)	Add according to manufacturer's recommendations.
Glyphosate products	Mix 1 part glyphosate with 4-9 parts water (10-20% solutions).
+ Surfactant	1 oz (2 tablespoons) per gallon of glyphosate mixture.
+ Ammonium sulfate	3 oz (6 tablespoons) per gallon of glyphosate mixture.
+ A marker dye (i.e., Blazon Blue)	Add according to manufacturer's recommendations.
Weedar 64	Mix 1 part Weedar to 2 parts water for hockey-stick application. Best results are obtained when used in late June and July. Do not drip or touch vines.

GLAUCOUS GREENBRIER *Smilax glauca* (Silverleaf sawbrier)**PRIORITY 1**

Greenbriers are very difficult to control because they have extensive underground storage organs. Glaucous greenbrier (silverleaf) is more difficult to control than common greenbrier (greenleaf). Glaucous grows in dense patches, spreads rapidly, and usually reaches just above vine level. It significantly impacts fruit production. Digging up the root system is not practical and will cause significant vine and bog damage. Infestations of sawbrier are more likely to occur on high edges or in locations where the bed is out of grade.

S. glauca may be managed by hand-wiping if sufficient coverage is obtained. Summertime wipes of Roundup may offer partial control. Clipping stems with Roundup-dispensing applicators in August may offer an additional partial control. Severe infestations of *S. glauca* may necessitate bog renovation. Weedar 64 may be used on sawbrier as per dewberry recommendations. Flooding is not effective against sawbrier.

WILD BEAN *Apios americana* (Ground Nut)**PRIORITY 1**

Wild bean can be partially controlled with both pre- and postemergence control techniques. Nuts may be dug out and removed from the bog. When using wipes, stake the wild bean vines to make wiping easier.

WILD BEAN - CHEMICAL RECOMMENDATIONS

Stinger	<i>Spray:</i> 0.07-0.11 oz/gal	0.4-0.7 tsp (1.8-3.0 ml) per gallon. Wild bean is very sensitive to Stinger.
	<i>Wipe:</i> 2.5 oz/gal	wipe: 5 TBSP per gallon (2% solution). See Notes on Stinger.
Casoron 4G	75 lb/A	Apply as two separate applications. Time applications before rain or irrigate within one day. Apply late March to early April. For best results, allow 10 days between applications.
FOLLOWED BY		
Devrinol 10G	75 lb/A or	
Devrinol 50DF	15 lb/A	
Salt	1 lb/gal water	Apply during the summer months after berry set. Never spray more than 200 gal/A. This practice keeps bean tops burned off. Repeat as necessary. Wash equipment with soap and water immediately after use.
Glyphosate products/ Roundup WeatherMAX		Apply anytime weeds are present except 30 days before harvest. For rates, see below and Notes on Roundup.

POISON IVY *Toxicodendron radicans***PRIORITY 1**

This particular weed seems to be getting worse on many production areas. Treat small infestations early and eradicate! Control is very difficult because the plant grows close to the vines. Glyphosate wipes will control this weed, but the potential for vine damage is high. Use of specialized applicators that minimize drip is recommended. Recent research has shown that clipping the stems with Roundup-dispensing applicators in August may offer partial control. In addition, research indicated that early-mid September applications may give better control than August applications. Again, this technique can be used on bogs that will not be harvested due to crop-destruct floods (or

24 Weeds

very late harvested bogs). Rates as low as 5-10% solutions gave decent control. Control for significant infestations of poison ivy is post-harvest spot treatments with Roundup sprays or mechanical spot renovation followed by replanting. Adding 1 part Weedar 64 to 4 parts of the Roundup mixture may be helpful but will likely increase the possibility of crop injury. See 2,4-D cautions.

Many people are highly allergic to poison ivy. Protective lotions and soaps are available that minimize the irritation caused by the poison ivy oils. Lotions are usually applied prior to exposure and soaps are used to remove oils after exposure to the plant. Rinse with a lot of cool water; small amounts of water may only spread the oils.

COMMON GREENBRIER *Smilax rotundifolia* (Greenleaf sawbrier, bullbrier) **PRIORITY 2**
Common greenbrier is bushier and spreads more slowly than glaucous greenbrier and grows well above the cranberry vines. It is easier to control with glyphosate wipes than glaucous greenbrier. Thorough coverage is important. Repeat applications in successive years may be needed for total control. See Notes on Roundup. Digging up the root system is not practical and will cause significant vine and bog damage.

YELLOW LOOSESTRIFE *Lysimachia terrestris* **PRIORITY 2**
Yellow loosestrife can cause moderate yield reductions. It may also serve as an early-season host for dodder and tips may harbor *Sparganothis* larvae. Preemergence herbicide application should be made in the spring. Loosestrife may be wiped with glyphosate during the season. Fall applications of Casoron may also offer some control of loosestrife. Since loosestrife is difficult to control, efforts should begin while patches are still small and before they have a chance to spread.

<u>CHEMICAL RECOMMENDATIONS</u>		
Casoron 4G	up to 100 lb/A	Spring or fall application. Offers partial control. See Notes on Casoron.
Roundup WeatherMAX / Glyphosate products		Apply anytime weeds are present except 30 days before harvest. See Notes on Roundup.
Weedar 64		Mix 1 part Weedar to two parts water for hockey-stick application. Best results are obtained when used in late June and July. Do not drip or touch vines.

NARROW-LEAVED GOLDENROD *Euthamia tenuifolia* **PRIORITY 2**
Narrow-leaved goldenrod causes little direct yield loss, but is extremely difficult to control and spreads rapidly. For this reason, every effort should be made to control patches before they spread. Preemergence and postemergence herbicides can provide partial control of this weed. Clipping the stems with Roundup-dispensing applicators in August may offer partial control.

<u>CHEMICAL RECOMMENDATIONS</u>		
Roundup WeatherMAX / Glyphosate products		Apply anytime weeds are present except 30 days before harvest. See Notes on Roundup.
Weedar 64		Mix 1 part Weedar to two parts water for hockey-stick application. Best results are obtained when used in late June and July. Do not drip or touch vines.
Stinger	Spray: 0.33-0.5 oz/gal Wipe: 2.5 oz/gal	0.75-1.2 tsp (3.6-5.4 ml) per gallon. See Notes on Stinger. wipe: 5 TBSP per gallon (2% solution).

ASTERS *Aster sp.* **PRIORITY 2**
The impact of asters on cranberries is variable. Asters are usually found in bare patches on the bog. Once the patch becomes established, asters are much harder to control. Applications of Casoron in March-April or November offer some suppression but will probably not eradicate this weed. Higher rates of Stinger offer the best option. Glyphosate wipes in the summer may be helpful. Thorough coverage and repeat applications are necessary.

CHEMICAL RECOMMENDATIONS FOR ASTERS

Casoron 4G	up to 100 lb/A	Apply Spring or Fall. See Notes on Casoron.
Iron sulfate	3 oz/sq. ft	Apply during Summer. See Notes on Iron Sulfate.
Weedar 64		Mix 1 part Weedar to two parts water for hockey-stick application. Best results are obtained when used in late June and July. Do not drip or touch vines.
Stinger	<i>Spray:</i> 0.33-0.5 oz/gal <i>Wipe:</i> 2.5 oz/gal	0.75-1.2 tsp (3.6-5.4 ml) per gallon. See Notes on Stinger. wipe: 5 TBSP per gallon (2% solution).

UPRIGHT BRAMBLE *Rubus allegheniensis* (Blackberry)**PRIORITY 2**

Upright bramble is not as common as the dewberries. It does not trail on the ground. Control may be obtained with glyphosate wipes. Young plants can be pulled or dug out by the roots. Weedar 64 may be used as described for other *Rubus* (dewberries) species.

CHOKEBERRY *Pyrus melanocarpa***PRIORITY 3**

Infestations of chokeberry can reduce yields and will spread in the beds. The best management strategy is to treat patches before they get too large. Chokeberry plants do not grow tall on the bog. Take extra care when using postemergence wipes of glyphosate products to minimize vine injury. When chokeberry plants are short, it may be more effective to hand-wipe them rather than wiping with a hockey stick wiper.

SEDGES**PRIORITY 3**

Management of sedges combines cultural and chemical controls. Hand dig, pull small patches or spot-treat with one of the preemergence herbicides listed below. Encourage vine growth in the bare areas so the sedges will not recolonize. Refer to the table below to locate the target weed and recommended control options and related information. TIMING: Spring applications are typically done from March through mid-April; Fall applications are typically done 1-2 weeks after harvest but at least 2-3 weeks prior to the winter flood. (S) or (F) following the weed name indicates 'Spring only' or 'Fall only' applications are preferred.

SEDGES - CHEMICAL RECOMMENDATIONS			
HERBICIDE	RATE	WEEDS CONTROLLED	NOTES
Devrinol 10G	60-90 lb/A	Nut sedge (nutgrass)	Rate for peat-based bogs.
Devrinol 50DF	40-60 lb/A		Rate for mineral soils.
	12-18 lb/A (peat)		Works best when applied to a weed-free surface.
	8-12 lb/A (mineral)		
Casoron 4G	up to 100 lb/A	Cottongrass, Dulichium(S), Fresh meadowgrass (F), Needlegrass, Nut sedge, Spike rush, Woolgrass	
Evital 5G	80-120 lb/A	Needlegrass (S), Nut sedge	Needlegrass is difficult to hand-pull.
	120-160 lb/A	Broomsedge, Needlegrass (F), Spike rush, Woolgrass	Broomsedge may be controlled with glyphosate wipes. Dig up clumps.
Weedar 64		Three-square	Mix 1 part Weedar to two parts water for hockey-stick application. Best results when used in late June and July. Do not drip or touch vines.

SHEEP LAUREL *Kalmia angustifolia***PRIORITY 3**

Sheep laurel can spread on the bog and reduce yields. Saplings should be pulled by hand. Larger plants can be wiped with glyphosate products (see Notes on Roundup).

26 Weeds

LEATHERLEAF *Chamaedaphne calyculata*

PRIORITY 3

Leatherleaf is a perennial, woody plant that can spread on the bog and reduce yield. It can be controlled by hand-wiping during the summer with a solution of glyphosate products. Weedar 64 can also be used as a wipe as per dewberry recommendations.

RUSHES *Juncus sp.*

PRIORITY 3

Rushes grow in clumps and can become quite large when well established. Control of large plants with preemergence herbicides may be difficult except at very high rates. Control may also be possible with hand-digging or repeated hand-wiping with glyphosate solutions. **TIMING:** Spring applications are typically done from March through mid-April. (S) following the weed name indicates if 'Spring only' applications are preferred. Use glyphosate wipes in summer.

RUSHES - CHEMICAL RECOMMENDATIONS

<u>HERBICIDE</u>	<u>RATE</u>	<u>WEEDS CONTROLLED</u>	<u>NOTES</u>
Devrinol 10G	60-90 lb/A 40-60 lb/A	Soft rush (<i>J. effusus</i>)	Rate for peat-based bogs. Rate for mineral bogs.
Devrinol 50DF	12-18 lb/A (peat) 8-12 lb/A (mineral)		Works best when applied to a weed-free surface.
Casoron 4G	up to 100 lb/A	Canada (S), Mud, Soft (S)	Offers partial control for Canada rush.
Evital 5G	120-160 lb/A	Canada rush	
Salt	1-3 teaspoons		Apply sodium salt to the base of each rush clump in the spring, prior to bud break.

PERENNIAL GRASSES

PRIORITY 3

These grasses often colonize bare areas. Encouraging vine growth may reduce potential for problems. Some species may be difficult to eradicate once established. Postemergence herbicides are now available for grass control on bearing and non-bearing beds. **TIMING:** Spring applications are typically done from March through mid-April; Fall applications are typically done 1-2 weeks after harvest but at least 2-3 weeks prior to the winter flood. (S) or (F) following the weed name indicates if 'Spring only' or 'Fall only' applications are preferred. Apply Devrinol early in spring (usually prior to April 10) for cutgrass control; it comes up earlier than other grasses.

CHEMICAL RECOMMENDATIONS – PREEMERGENCE OPTIONS FOR PERENNIAL GRASSES

<u>HERBICIDE</u>	<u>RATE</u>	<u>WEEDS CONTROLLED</u>	<u>NOTES</u>
Devrinol 10G / 50DF	60-90 lb/A / 12-18 lb/A 40-60 lb/A / 8-12 lb/A	Poverty grass Rice cutgrass	Rate for peat-based bogs. Rate for mineral soils. Repeated mowing is helpful.
		Summergrass (S)	Early application is desirable (before April 10) for best control. Improve drainage for cutgrass problems. Hard to mow.
Casoron 4G	up to 100 lb/A	Mannagrass (S) Rattlesnakegrass (S)	Clean ditches in infested areas. Minimal control by pulling.
		Rice cutgrass, Summergrass, Velvetgrass	
Evital 5G	80-120 lb/A 120-160 lb/A	Rice cutgrass (S), Smokegrass (S) Broad-leaved panicgrass (S) Summergrass Switchgrass (F)	Gives partial control. Hard to mow. Difficult to control. Repeated mowing helps. Best to dig out plants prior to seed formation.

PERENNIAL GRASSES - CHEMICAL RECOMMENDATIONS - POSTEMERGENCE OPTIONS

Poast	1-1.5% solution + Dash HC (0.5%) OR crop oil concentrate (1%)	True grasses only. Apply at 6-8 leaf stage for best results.
Mix 2 oz Poast + 0.6 oz Dash HC or 1.3 oz crop oil concentrate per gallon. Repeat applications may be needed for these grasses. Addition of other adjuvants is not recommended. May be applied by broadcast applicator or by air. Chemigation is not permitted! See Notes on Poast.		
Select	6- 8 oz/A per application + 1% v/v COC in the finished spray volume unless label indicates otherwise.	True grasses only. Chemigation not permitted. Repeat applications may be needed. Use 10-30 gal water per acre. For each gal, mix 0.5 oz (1 TBSP) Select + 1.3 oz COC (2.5 TBSP) for a mid-range rate. Do not apply between hook and full fruit set.
Select MAX	9-16 oz/A per application + 0.25% v:v non-ionic surfactant (NIS)	See Select. Use 1.3 TBSP (39 ml) Select Max + 0.65 TBSP (19 ml) NIS per gallon for a mid-range rate. Use NIS (not COC) with Select Max.

RED MAPLE and other trees**PRIORITY 3**

The best management strategy is to pull young saplings before the root system becomes established (hopefully less than 1 year old). Larger trees must be dug out. Glyphosate wipes may be used to control small maples and to weaken large trees to facilitate removal. Clipping stems with Roundup-dispensing applicators in August may offer partial control.

CHEMICAL RECOMMENDATIONS**Roundup WeatherMAX /
Glyphosate products
Weedar 64**

Apply anytime weeds are present except 30 days before harvest. See Notes on Roundup. Mix 1 part Weedar to two parts water for hockey-stick application. Best results are obtained when used in late June and July. Do not drip or touch vines. Labeled for red maple only.

MOSESSES [*Haircap moss (Polytrichum sp.)*, *Sphagnum moss (Sphagnum sp.)*]**PRIORITY 4**

The presence of these plants on the bog may indicate a drainage problem. Evaluate the drainage in the area and improve prior to starting a chemical control plan.

HAIRCAP MOSS - CHEMICAL RECOMMENDATIONS

Casoron 4G	up to 100 lb/A	See Notes on Casoron. Be judicious in applying high rates for moss control, especially on vines showing signs of stress.
Ammonium Sulfate	15 oz/100 sq. ft	21-0-0. Apply in the spring.

SPHAGNUM MOSS - CHEMICAL RECOMMENDATIONS

Iron sulfate	3 oz/sq. ft	Apply in March to mid-April. See Notes on Iron Sulfate.
Casoron 4G	up to 100 lb/A	Apply in the fall only. See Notes on Casoron.

JOE-PYE WEED *Eupatorium dubium***PRIORITY 4**

This perennial plant generally grows along the ditch areas. Due to its tough stem, it may be difficult to hand-weed. Some growers use pliers to uproot large plants. Glyphosate wipes during the summer offer good control. Stinger should also control this weed.

28 Weeds

CINQUEFOIL *Potentilla canadensis, P. simplex* (Five-finger)

PRIORITY 4

The impact of cinquefoil is variable. Colonization of cinquefoil may indicate a problem with vine growth and high pH (too basic). Improvement of fertilizer program may help control this weed. Hand-wiping or hand-pulling can also be used to eliminate small patches. If the soil is considered alkaline, the use of sulfur may help improve the soil condition such that the vines become more competitive. See Notes on Sulfur.

CHEMICAL RECOMMENDATIONS FOR CINQUEFOIL

Iron sulfate 20%	3 oz/sq. ft	Apply during the summer months. Several formulations and percent active ingredient of iron sulfate are available. Granular forms are easier to apply (drop-spreader), but take longer to act than finely powdered formulations. See Notes on Iron sulfate.
-------------------------	-------------	--

MEADOWSWEET *Spiraea latifolia*

PRIORITY 4

Meadowsweet is a slow spreader on cranberry bogs. This weed should be pulled out by hand or wiped during the summer.

CHEMICAL RECOMMENDATIONS

Roundup WeatherMAX / Glyphosate products	Apply anytime weeds are present except 30 days before harvest. See Notes on Roundup.
---	---

WHITE VIOLET *Viola lanceolata*

PRIORITY 4

Found most often in bare patches on beds, white violet is thought to compete poorly with established vines. Thus, the best management approach is the fill in bare spots on the bog by encouraging runner growth. Otherwise, hand-pulling is the next best recommendation, especially on new bogs. No chemical controls are recommended.

PITCHFORK, RAGWEED, AND FIREWEED (A)

PRIORITY 4

Bidens frondosa, Ambrosia artemisiifolia, and Erichthites hieracifolia (Beggarstick, stick-tights, Spanish needles/Common ragweed, hogweed, bitterweed, Roman wormwood/Cottonweed, pilewort.)

PITCHFORK, RAGWEED, & FIREWEED - CHEMICAL RECOMMENDATIONS

Casoron 4G	up to 100 lb/A	Apply in spring. May also be applied in the fall for control of ragweed. Moderate rates are suggested.
Caliber 90	up to 3.3 lb/A	Apply in spring. See Notes on Princep.
Princep 4L	up to 3 qt/A	
Princep 80W	3.75 lb/A	
Weedar 64	for ragweed only	Mix 1 part Weedar to two parts water for hockey-stick application. Best results are obtained when used in late June and July. Do not drip or touch vines.
Stinger	<i>Spray:</i> 0.07-0.11 oz/gal <i>Wipe:</i> 2.5 oz/gal	0.4-0.7 tsp (1.8-3.0 ml) per gallon. Pitchfork and ragweed are sensitive to Stinger. Wipe: 5 TBSP / gal (2% solution). See Notes on Stinger.

HARDHACK *Spiraea tomentosa* (Steeplebush)

PRIORITY 4

Hardhack is a solitary plant that does not spread except through seed. Wiping or pulling this weed is very effective, but the roots of hardhack can be woody and larger plants may cause damage to the bog when pulled.

CHEMICAL RECOMMENDATIONS

Roundup WeatherMAX/ Glyphosate products	Apply anytime weeds are present except 30 days before harvest. See Notes on Roundup.
--	---

ANNUAL GRASSES**PRIORITY 4**

Generally, annual grasses are most likely to appear in bare patches or on new bogs. Hand-pulling and/or treatment with postemergence grass herbicides may be used on new and established bogs. Encourage vine growth to reduce ability of weeds to colonize the bog surface. Preventing seed production may also be important in managing these weeds. **TIMING:** Spring applications are typically done from March through mid-April. (S) following the weed name indicates if 'Spring only' applications are preferred.

CHEMICAL RECOMMENDATIONS FOR ANNUAL GRASSES - PREEMERGENCE OPTIONS

HERBICIDE	RATE	WEEDS CONTROLLED	NOTES
Devrinol 10G / 50DF	60-90 lb/A / 12-18 lb/A 40-60 lb/A / 8-12 lb/A	Barnyardgrass, Corn grass, Warty panicgrass (S)	Rate for peat-based bogs. Rate for mineral soils. Take action to promote vine growth.
Casoron 4G	up to 100 lb/A	Crabgrass (S)	Take action to promote vine growth.
Evital 5G	120-160 lb/A	Barnyardgrass, Corn grass	Take action to promote vine growth.

Lower rates have been shown to be effective on established bogs. Use the minimum effective dose. Very low rates (<25 lb/A) have caused injury on newly planted vines.

ANNUAL GRASSES - POSTEMERGENCE OPTIONS

Poast	1-1.5% solution +Dash HC (0.5%) OR crop oil concentrate (1%)	True grasses only!	For each gallon, mix 2 oz Poast plus 0.6 oz. Dash HC or 1.3 oz crop oil concentrate. See Notes on Poast.
Select	6-8 oz/A per application +1% v/v crop oil concentrate (COC) in the finished spray volume unless label indicates otherwise.		True grasses only. Chemigation not permitted. Repeat applications may be needed. Use 10-30 gal water per acre. For each gal, mix 0.5 oz (1 TBSP) Select + 1.3 oz COC (2.5 TBSP) for mid-range rate. Do not apply between hook and full fruit set.
Select Max	9-16 oz/A per application + 0.25% v:v non-ionic surfactant (NIS)		See Select. Use 1.3 TBSP (39 ml) Select Max 0.65 TBSP (19 ml) NIS per gallon for a mid- range rate. Use NIS (not COC) with Select Max.

CLOVER AND VETCH *Trifolium repens, and Vicia sp.***PRIORITY 4**

Clover and vetch tend to occur in areas of very high pH (alkaline soils). If soil pH is 5.0 or above, spot treat with two applications of sulfur at the rate of 0.2 oz/sq. ft. Apply in the late spring when soil is drained and frost protecting is over. For more details, refer to Notes on Sulfur. Lower rates of herbicides may be effective when sulfur has been added and the pH is lowered. **Stinger** offers good postemergence control of these weeds (use rates similar to asters and narrow-leaved goldenrod). See 'Stinger notes' at the end of the weed section. Control clover early spring prior to budbreak.

FEATHER, ROYAL, CINNAMON, AND SENSITIVE FERNS**PRIORITY 4***Dryopteris thelypteris, Osmunda regalis, Osmunda cinnamomea, Onoclea sensibilis*

(S) or (Smr) following the weed name indicates if "Spring" or "Summer" applications are preferred. When using iron sulfate treatments, apply a small amount to each plant. See Notes on Casoron and Iron Sulfate.

30 Weeds

<u>FERNS - CHEMICAL RECOMMENDATIONS</u>			
<u>HERBICIDE</u>	<u>RATE</u>	<u>WEEDS CONTROLLED</u>	<u>NOTES</u>
Casoron 4G	up to 100 lb/A	Bracken fern (S) Royal fern (S)	Appears on bogs showing signs of stress. Spot-treat and use moderate rates. Spot-treat and use moderate rates.
Iron sulfate	2 oz/sq. ft [20% a.i.] See Iron Sulfate notes	Feather fern (Smr) Sensitive Fern (Smr)	Sensitive fern is difficult to hand weed due to perennial rhizomes breaking. Use caution on bogs that have been sanded within 18 months.
Iron Sulfate & Salt	9:1 ratio (iron:salt)	Cinnamon fern, Feather fern, Sensitive fern	Treat during the summer months. Place a small amount at the base of each plant.

MINOR WEEDS

PRIORITY 4

[Blue joint (*Calamagrostis canadensis*), Hawkweed (*Hieracium sp.*), Horsetail (*Equisetum arvense*), Marsh St. John's Wort (*Triadenum sp.*), Orange-grass/pinweed (*Hypericum gentianoides*), *Hypericum* species, Sweet pepperbush (*Clethra alnifolia*), Plantain (*Plantago* or *Littorella sp.*), Smartweed (*Polygonum sp.*), Sorrel (*Rumex sp.*), and Wild Strawberry (*Fragaria sp.*)]

If weeds are scattered or of minor importance, consider potential vine stress or injury when choosing herbicide and rate. Consider hand-pulling instead. For localized weed patches, consider spot treatment. Postemergence treatment with glyphosate products may also offer some control. Combine herbicide treatments with steps to improve vine growth. Control of *Equisetum* generally requires more than 75 lbs/A Casoron. This weed is often confused with orange-grass; the same common name is often used for both plants in MA but orange-grass is much easier to control.

<u>CHEMICAL RECOMMENDATIONS</u>	
Casoron 4G	Use lowest rates possible. Apply in spring or fall. See 'Notes on Casoron'.
Weedar 64	Mix 1 part Weedar to two parts water for hockey-stick application. Best results are obtained when used in late June and July. Do not drip or touch vines. Labeled for <i>Hypericum</i> species and sweet pepper bush only.

ALGAE (*Green scum*) Algaecides are usually prescribed on an acre-foot basis. An acre-foot is the amount of water needed to cover one acre of bog with one foot of water (~300,000 gallons of water, assuming the bog is level).

<u>ALGAE - CHEMICAL RECOMMENDATIONS</u>		
Copper sulfate	4 lb/ acre-ft	Apply evenly on ice or in bog waters. When bogs are treated during winter months, water should be impounded for 5-7 days to allow for degradation in cool-temperature water. May also be used in late water.
Algae-Pro	0.75-1.5 gal/A-ft	Amount will vary depending on product, water volume, and algal density. Carefully read the label before application. See Notes on
Cutrine-Plus	0.6-1.2 gal/A-ft	Copper Sulfate and Copper Complexes.

OUTSIDE OF BOG AREA – WEED MANAGEMENT

<u>AQUATIC WEEDS -</u>		<u>CHEMICAL RECOMMENDATIONS</u>
Diquat Reward	1-2 gal per surface acre	Use during summer months. Water use is restricted for various time periods depending on product and pattern of use. CHECK THE LABEL! Use only on <u>still water</u> areas outside of bog (i.e., farm ponds, reservoirs). DO NOT USE IN OR ON BOG DITCHES.
Rodeo	1.25% solution	Apply during the summer months. Rodeo is registered for use on noncrop land only. Use in interior ditches is not permissible. Recommended spray solution: 5 oz/3 gallons. <u>Add a nonionic surfactant</u> at the rate of 0.25-0.50% volume basis (1-2 oz or 2-4 Tbsp in 3 gal.).

WOODY AND BROADLEAF PERENNIALS (not in ditches or canal banks)

Hand-pulling is most beneficial in the spring and early summer when the soil is moist and the plants are fairly small.

CHEMICAL RECOMMENDATIONS – WOODY PERENNIALS – NON-BOG USE ONLY!!

Weedone CB	Do not dilute.	Apply in February and March. Spray to wet. Avoid drift onto bog. Controls woody plants on roadsides and non-crop areas. It is no longer produced, but available product may be used <u>off the bog</u> .
Crossbow	up to 2 gal/A	Mix with enough water to deliver 10-30 gal/A. Application rates vary depending on target species and application method. Drift to desirable plants may cause injury (esp. grapes and tomatoes). Do not apply to water. Read the label! See Notes on 2,4-D.

DITCH MANAGEMENT

WOODY AND BROADLEAF PERENNIALS ON DIKES (BOG-SIDE).

Cultural controls include mowing the ditch and dike areas during the summer months. Some areas may need to be done more than once. Hand-pulling is most beneficial in the spring and early summer when the soil is moist and the plants are fairly small. Controlling weeds on the dikes may be useful in reducing spread of these weeds onto bogs.

DITCH WEEDS (*i.e., Arrowhead, Pickerelweed, Pond lilies, Bur-reed, Duckweed*)

Clean ditches by hand or mechanically preferably twice a year. Draining ditches can sometimes be helpful in killing some aquatic weeds (*i.e., duckweed*). Preemergence herbicides registered for use on the bog may **NOT** be used in the ditches for weed control.

DITCH WEEDS - CHEMICAL RECOMMENDATIONS

Roundup WeatherMAX	Use as a <i>wipe</i> during the summer months in dry ditches. See Notes on Roundup.
Roundup WeatherMAX	Use as a <i>spray</i> during the summer months in dry ditches. Use a 1-1.5% solution on a volume-to-volume dilution. Spray to just wet vegetation, not to run-off. Ditches must be kept dry at least 2 days after application.

32 Weeds

CAUTIONS AND OTHER NOTES

1. Chemicals not registered for use on cranberries must not be used.
2. Herbicide use may weaken vines and crops may be reduced.
3. To be most effective, rain should follow the application of any dry herbicide formulation within 4 days or the bog should be irrigated.
4. Wash equipment with soap (or detergent) and water immediately after using. Rinse with ammonia after using hormone-type herbicides (such as 2,4-D).
5. Hand-wiping with glyphosate products is often practical with some weeds if roots are weakened. This is particularly useful for dewberries after late water or a summer flood.
6. Mowing of tall weeds helps to prevent shading and reduces seed formation.
7. Late water causes general reduction of annual grasses and may reduce dewberry populations and re-growth.
8. Agricultural burning of brush or grass is allowed under regulations from the Director of Air Pollution Control, Southeastern Office of the Dept. of Environmental Protection and under permit from the local fire chief.
9. Review the Weed Management BMP in the UMass Best Management Practices Guide.

NOTES ON THE USE OF COMMON HERBICIDES

CASORON (Dichlobenil). Applications of Casoron are most effective when applied as close to the time of weed germination or emergence as possible. Since Casoron volatilizes quickly, it must be washed in by irrigation or rainfall ASAP after application. Apply pre-budbreak or post-harvest. Avoid applying during warm temperatures. Application just prior to sanding or on weak or new vines may cause injury. Applications on top of sand or late applications can be made, but must be watered in *immediately*. Low rates (<40 lb/A) may be applied after removal of a late water flood to control dodder with minimal risk of phytotoxicity. In general, applications of preemergence herbicides are not recommended *after* the later water flood is withdrawn.

Casoron may be applied by air or by ground equipment. Multiple applications may be made as needed. Allow an interval of 3-6 wks between applications. Do not exceed 100 lb/A in any 12-month period. Single doses of high rates of Casoron may be needed to control some perennial weeds. However, some weeds are not controlled by Casoron at any rate due to their deep root systems. Cranberry vines with weakened root systems are more susceptible to stresses such as drought and may become more stressed with herbicide application. Some vine injury may occur from herbicide applications made in areas where puddling is a problem.

Distribute Casoron uniformly. Avoid overlapping of herbicide. Temporary reddening of vines may occur especially with late spring application or when applied on sandy bogs. Do not apply after bud elongation as vine injury will occur and yields may be reduced. Do not apply to young beds (less than 3 yrs old unless root systems are well established) or on bogs prior to or immediately after mowing vines. Do not sand (spring or fall) on top of a Casoron application. Casoron is labeled for application in the fall prior to ice sanding that winter or in the spring after ice sanding. The efficacy of fall applications for many weeds has not been documented, but growers have reported good success in some cases.

COPPER SULFATE and COPPER COMPLEXES (Algae-Pro, Cutrine-Plus). Copper sulfate may be used to control algal growth on winter or late water floods. Cutrine-Plus and Algae-Pro work best when water temperatures are warm (~ 60°F). These copper-complex products are formulated to last longer than copper sulfate in hard water (carbonates present). Copper-complex products work best when applied under calm and sunny conditions.

If you are holding a 4-week late water flood, plan to apply a copper product mid-way through the flooding cycle (for more details, see Prevention of Scum in the Late Water section). These products are typically applied directly through the irrigation systems (with heads on risers). For winter floods or late water floods of short duration, scout for algae and apply when growth is first visible on the water surface. Remember, these products only prevent further algal growth; they do not kill or remove what has already grown. Crystal copper sulfate will dissolve easily in water and can be applied as a spray solution. Though somewhat uncommon now, crystalline products can be placed in burlap

bags and dragged across the water surface. When bogs are treated with copper sulfate during the winter months, water should be impounded for one week. Since late water floods tend to be warm, there is no need to impound these waters. Do not apply to water except as directed on the label. These products are toxic to fish.

CROSSBOW, WEEDAR 64, WEEDONE CB (2,4-D). Crossbow and Weedone CB are labeled for **non-bog use only**. Be cautious! Crossbow contains trichlopyr for which there is **no food tolerance**. Do not use Crossbow or Weedone CB on dikes or canal banks. Use it only on weeds located far away from the bog. These products have considerable potential to evaporate and cause crop injury. Avoid applying 2,4-D on hot, sunny, and humid days when there is little air movement. Weedar 64 is the only 2,4-D product that can be used on the bog. 2,4-D products can be highly effective at controlling some weeds. However, the potential for significant vine injury may outweigh the advantages of using these materials. Please Note: Weedar 64 now has a 30-day PHI and may only be applied once per season.

DEVIRINOL (Napropamide). Devrinol 50DF is now available (as a 24c label) for use on cranberries. Unlike Devrinol 10G, the new formulation can be injected through the irrigation system. We have some experience with 50DF and it seems to perform at least as well as Devrinol 10G. Growers who used it in 2005-06 reported good control. Be sure to get the supplemental label at the point of purchase. Since Devrinol will control grasses, give careful consideration to use of 50DF if you are establishing ditch banks and your irrigation system delivers coverage to the ditch bank area.

Do not exceed 30 lb/A Devrinol 10G or 6 lb/A Devrinol 50DF on new plantings. On established bogs, this herbicide provides some control of grassy weeds and annual broadleaf plants at higher rates (10G: 60-90 lb/A; 50DF: 12-18 lb/A for peat beds; 10G: 40-60 lb/A; 8-12 lb/A 50DF for mineral soils), but works best on weed-free areas. For best results, water in immediately. 10G may be applied by air or ground; 50DF can be chemigated. Using the appropriate rate for the age of the bog. Devrinol can be used under or on top of sand.

DIQUAT. This herbicide should only be used on water weeds growing in areas OUTSIDE OF THE BOG. Do not use in any ditch associated with the production area. Diquat will control water weeds such as bladderwort, coontail, elodea, and pondweeds. A non-ionic surfactant (e.g. X-77) may improve performance. Check the label for rate information.

EVITAL (Norflurazon). Vine injury may occur in areas where water stands several days after flooding or heavy rains. **Be very conservative when applying Evital to new plantings!** Vines have shown severe phytotoxicity to rates as low as 25 lb/A when applied 3-4 weeks after planting. Do not apply more than 80 lb/A on newly planted bogs. Do not apply more than 160 lb/A per season on an established bog. Use lower rates on stressed vines or sensitive cultivars such as Stevens and McFarlin. Growers have reported good results with low rates (50-75 lb/A) for fall applications on these varieties; spring applications should not exceed 60 lb/A. Sanding can be done on top of an Evital application, but it is not recommended especially on bogs that have drainage problems. Sanding after applications of 50 lb/A or less has given good results. Growers have reported that applications of Evital (50-60 lb or less) on top of sanded vines work adequately on healthy well-drained beds.

FUMIGANTS. Basamid (dazomet) and Vapam (metam-sodium) are soil fumigants that can be used on beds scheduled for renovation. **DO NOT USE FUMIGANTS AS A SPOT-TREATMENT IF ANY VINES WITHIN A DIKED SECTION WILL BE HARVESTED.** If you are renovating an entire section, a portion of that section can be spot-treated with a fumigant. See 'New Plantings' section for more information on the use of fumigants.

IRON SULFATE. May be spread as a broadcast application through conventional fertilizer rigs, such as hand cranks. Traditional use has been with a 20% ferrous sulfate (fine powder) product, but other formulations are available. Application rates are listed in the Chart Book for the 20% a.i. product. Adjust accordingly if using another percent active ingredient. Iron sulfate at rates exceeding 1.1 oz/sq. ft (20% a.i. product) may kill vines if they have been sanded within the past 18 months. Do not use on new bogs. To be most effective, rain should follow within 4 days of an iron sulfate application or the bog should be irrigated. When a 9:1 iron sulfate to salt combination is used, rain or sprinkling is not necessary.

34 Weeds

POAST (Sethoxydim). This herbicide effectively controls emerged annual and some perennial *true grasses*. Sedges are not controlled. It may be used on bearing and non-bearing beds. There is a 60-day PHI on bearing beds. Do not apply more than 5 pints (80 oz) of product per season. Allow a minimum of 14 days between repeat applications. Phytotoxicity may result if the herbicide is applied during the heat of the day or during bloom. Application during cool periods of the day, but after dew has dried, is preferable. If you have never used this material before, it may be wise to test a small area before applying the product to a larger area.

Efficacy is enhanced by addition of crop oil concentrate (COC) or Dash HC, and either adjuvant should always be used. Other adjuvants may reduce efficacy or increase crop injury. Since COC or Dash HC can be mildly phytotoxic, Poast should not be applied during periods of crop stress or during flowering. Poast should not be mixed with other chemicals, particularly chemicals whose label warns against inclusion of an adjuvant.

Poast must be absorbed into the grass to be effective. Therefore, do not apply Poast if rainfall or irrigation is expected within one hour of application. Poast should be applied when grasses have 6 to 8 leaves to provide enough leaf surface for absorption. Apply Poast to grasses that are actively growing and free of stresses such as drought, disease, or mechanical injury.

POAST CANNOT BE APPLIED THROUGH THE IRRIGATION SYSTEM! Spot treatments with small sprayers are effective. **For one gallon Poast solution, mix 2 oz of Poast with 0.6 oz of Dash HC (or 1.3 oz of crop oil concentrate) in 1 gallon water.** Thoroughly wet the grass foliage, but do not let the solution run off the leaves.

Broadcast Application. Use standard high-pressure hollow cone or flat fan nozzles only. Use 5-20 gal of spray solution per acre at 40-60 psi. Inadequate coverage of grasses due to heavy cranberry canopy may reduce control. Do not use re-circulating sprays, wiper applicators or shielded applicators. Use of Poast with control drop application is not recommended due to erratic coverage. *Aerial Application.* Do not apply if wind speed is greater than 10 mph.

PRINCEP (Simazine). The Caliber 90 and Princep 4L formulations must be sprayed evenly with continuous agitation. Application through the sprinkler systems is not allowed. Apply before budbreak or in the fall after harvest. High rates may injure vines or crop. Thin or weak vines and new plantings one week to three years old are very susceptible to injury. Growers under Zone II regulations may not apply simazine (see Groundwater Protection and Zone II Section).

RODEO. This glyphosate product can only be used to control weeds that occur in dry ditches and canals outside of the production area. Application is spray to wet leaf surfaces, not to runoff. Extremely cool or cloudy weather following application may slow the activity of this herbicide. Best control is obtained when plants are at late growth stages approaching maturity. Weeds under stress will not be controlled as well as healthy plants. Rainfall within 6 hours of application may reduce effectiveness and heavy rainfall within 2 hours of application may necessitate reapplication. Do not add ammonium sulfate to Rodeo mixtures.

ROUNDUP (Glyphosate). Many RoundUp and glyphosate products are available on the market. Please read the label of any product you are using to ensure compliance. This product may be applied on bogs, by wiper or clipper, during the growing season. If you are using Round Ultra, you will need supplemental labels for dry ditch and postharvest sprays. If you are using WeatherMAX, these spray uses are incorporated into the label and additional labels are not needed. WeatherMAX is slightly more concentrated than Ultra, so keep that in mind when preparing solutions. Use 1%-1.5% solutions (2.5-3.8 TBSP or 38-57 ml/gal) for dry ditch applications and 0.4%-0.7% solutions (~3.0-5.5 tsp or 15-27 ml/gal) for postharvest sprays. Recent research indicates that Howes may be slightly more sensitive to postharvest spray injury than Early Black.

It is not necessary to mix Roundup Ultra or WeatherMAX with any additional surfactants or additives (as with older glyphosate products). Add a dye to track leaf coverage. Technical information indicates that ammonium sulfate may improve uptake of these Roundup products when moderate to large amounts of carbonates ('hard water') are present in water (rarely a problem in MA). Roundup WeatherMAX is rainfast 1-2 hours after application. Available glyphosate products vary as to whether they carry a 'Caution' label or 'Warning' label. Look at the label!! When using Roundup, protective eyewear is not mandated; the REI for WeatherMAX is 4 hr. Thorough coverage is essential to maximize control of perennial weeds. Do not touch or allow material to drip onto vines. **Apply any time weeds are present except 30 days before harvest.** Make herbicide mixtures fresh each day for maximum effectiveness. Do not store in galvanized containers.

Roundup Products	Mix 1 part glyphosate with 4-9 parts water (10-20% solutions). No additional additives, buffers, or surfactants are needed. However, the addition of ammonium sulfate may sometimes improve performance.
+ A marker dye (i.e., Blazon Blue)	Add according to manufacturer's recommendations.
Glyphosate products	Mix 1 part glyphosate with 4-9 parts water (10-20% solutions).
+ Surfactant	1 oz (2 tablespoons) per gallon of glyphosate mixture.
+ Ammonium sulfate	3 oz (6 tablespoons) per gallon of glyphosate mixture.
+ A marker dye (i.e., Blazon Blue)	Add according to manufacturer's recommendations.

OTHER GLYPHOSATE PRODUCTS. Glyphosate is sold under several product names. **CHECK THE LABEL!** To enhance control with glyphosate products other than Roundup WeatherMAX, add a nonionic surfactant (i.e., X-77) and ammonium sulfate (see rates above). Other label differences include: Do not apply if rainfall is expected within 6 hr of application. Do not irrigate within 6 hr of application. The REI is 12 hr for these products. Note also that glyphosate products other than Roundup WeatherMAX may carry a 'Warning' label, instead of a 'Caution' label. Always use a dye to track your coverage with any wipe product. Check the label for appropriate protective clothing.

Clipper Applicators (Roundup only). Concentrations of 50-100% Roundup have worked well. The herbicide should flow out consistently, but not so fast that herbicide drips from the blades. Be sure to use a dye. Clip weeds close to the ground, without contacting the vines. Roundup must contact the stem as you are cutting! 'Clip and dab' or 'mow and wipe' techniques may have reduced efficacy as the herbicide is not applied simultaneously with the cut. Late-season treatments give better results than early-season treatments. The effectiveness of post-harvest treatments with clippers is not known. Be sure to clean the blades after use to prevent corrosion. Availability of commercial clippers has become more scarce over the past few years. Growers may need to manufacture their own clippers.

General Wiping Tips. Use a small sponge or applicator that permits excellent coverage with minimal dripping. Adequate coverage of each stalk must be obtained. Several leaves (at least 50%) on each stalk must be treated with the herbicide. Repeat applications to remaining plants the following year. Be patient. Most treatments will not give 100% control in the first year. Applications in subsequent years should be less time-consuming.

Hand-wipe Technique for Controlling Dewberries or Other Weeds that Lay in the Vine Canopy. Application by hand with sponges or specially designed applicators may be necessary with low-growing weeds (e.g., bristly dewberry, poison ivy). Repeat applications within a season are legal and may be necessary, especially for well-established perennial weeds. Poor growing conditions such as drought stress, disease, or insect damage may reduce effectiveness. Avoid touching or dripping material onto cranberry plants during application. Some growers have had success staking the vines (tomato stakes or similar) for wiping and allowing them to dry prior to laying them back on the vines. This certainly reduces vine injury.

SUPPLEMENTAL LABEL USES. Supplemental labels are needed only if you are using Ultra for post-harvest sprays (0.5%-1%), applied as a spot-treatment, or sprays in dry ditches (1%-2%); these uses are permitted under regular labeling for WeatherMAX. These labels can be obtained at a local ag supplier or by request from the Cranberry Station. If you are unsure about using a supplemental label, call the Cranberry Station (ext. 21) for information.

SALT. Salt (sodium chloride) may be used as a spot-treatment for control of certain weeds (e.g., wild bean, rushes). Judicious applications do not inhibit re-colonization of cranberry vines once the weed dies. Do not use during bloom. Use of calcium chloride or other types of salts is not recommended. Salt is corrosive to machinery. Be sure to wash equipment thoroughly after application.

SELECT (clethodim). Similar to Poast with regards to target species, timings, and applicators. Note these differences. Apply when weeds are 2-6 inches high and actively growing (check label for specific heights for each target weed). Multiple applications are typically needed for perennial weeds. Results are best when weeds are not under stress. Irrigation should follow within 7 days if rainfall does not occur. Use of crop oil concentrate (COC) is recommended with Select; a non-ionic surfactant (NIS) can be used with Select Max. The potential for crop injury is much less with an NIS. For a 1-gal mixture, use 1 TBSP Select with 2.5 TBSP COC; use 1.3 TBSP Select MAX with 0.65 TBSP NIS. Select and Select MAX have a 30-day PHI. Do not apply more than 8 or 16 oz Select and Select

36 Weeds

MAX per application per acre, respectively. Do not exceed 32 or 64 oz per acre per season, respectively. Allow 14 days between applications. You can use 3-10 gallons water with aerial applications, 10-30 gal/A otherwise.

STINGER (clopyralid). Stinger is a selective, postemergence herbicide used to control: wild bean, narrow-leaved goldenrod (NLGR), asters, clover, ragweed, pitchfork (and other members of the Composite and Legume families), and certain other weeds within the treated area. Growers have reported effective control (and reduced vine injury) when using lower rates than recommended on the label. This is particularly true for wild bean control.

Apply when weeds are actively growing. It is best to apply Stinger when vines are dormant, if possible. For weeds that emerge late (NLGR, wild bean, etc.), wait until after fruit set to apply. It is not recommended to apply Stinger when vines are going through active growth spurts (e.g. budbreak-roughneck stage). Stinger has a 50-day PHI. Stinger may be applied as a wipe or as a spray. Spray to just wet the weeds, but not to run-off. BE VERY CAREFUL! Overspray can cause injury that may take 1-3 years for full vine recovery. Minimize drift when applying as a spray. Results may be slow to show; be patient. Two applications per season are permitted, not to exceed a total of 1 pint per acre. Stinger cannot be applied by air or through the irrigation system.

SULFUR. Determine soil pH in the weedy area prior to sulfur application. If pH is 5.0 or above, use two applications of 500 lb/A each (or 4 applications of 250 lb/A) to reach 1,000 lb. of elemental sulfur per season. Begin application in late spring when soil is drained and sprinkling for frost is over. Most growers allow 3-6 weeks between applications. Do not apply sulfur to puddled or waterlogged areas as resultant production of hydrogen sulfide can cause severe vine toxicity. Changes in pH can be very slow. Granular applications may take up to nine months to reduce pH enough to affect weed populations. The smaller the sulfur pellet size, the faster the pH is lowered. Use pelletized sulfurs only. Do not use flours of sulfur; they can be phytotoxic and are difficult to apply. Do not use potassium sulfate. Yearly sulfur applications may be needed as the pH can creep up in subsequent years. Test soil pH yearly to determine the effectiveness of sulfur applications. The effect of lowered pH on control of cinquefoil is moderate. Eye protection and dust masks are recommended when making sulfur applications.