



Active Ingredient:  
 [S]-*trans*-2-Amino-4-(2-aminoethoxy)-  
 3-butenic acid hydrochloride . . . . . 15%w/w  
 Other Ingredients . . . . . 85%w/w  
 Total . . . . . 100%

EPA Reg. No. 73049-45  
 EPA Est. No. 33762-IA-001

**FOR USE ON APPLE, CUCUMBER, PEAR,  
 PECAN, STONE FRUIT, AND WALNUT**

**INDEX:**

- 1.0 First Aid
- 2.0 Precautionary Statements
  - 2.1 Hazard to Humans and Domestic Animals
  - 2.2 Personal Protective Equipment (PPE)
  - 2.3 User Safety Recommendations
  - 2.4 Environmental Hazards
- 3.0 Directions for Use
- 4.0 Aricultural Use Requirements
- 5.0 General Information for Use
- 6.0 General Information Instructions
- 7.0 Compatibility with Other Agriculture Products
- 8.0 Apple and Pear
- 9.0 Stone Fruit
- 10.0 Nut, Pome and Stone Fruit Crops
- 11.0 Cucumber
- 12.0 Storage and Disposal
- 13.0 Notice to User

**KEEP OUT OF REACH OF CHILDREN  
 CAUTION**

1.0

<b>FIRST AID</b>	
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

(continued on next column)

<b>HOT LINE NUMBER</b>
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-892-0099 (24 hours) for emergency medical treatment and/or transport emergency information. For all other information, call 1-800-6-VALENT (682-5368).</p>

**2.0 PRECAUTIONARY STATEMENTS**

**2.1 HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
 CAUTION**

Harmful if inhaled or absorbed through skin. Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

**2.2 Personal Protective Equipment (PPE)**

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Waterproof gloves

Hand-wand applicators must wear dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with N, R, P or HE filter when applying this product.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**2.3 User Safety Recommendations**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then, wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**2.4 Environmental Hazards**

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

**3.0 DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State/Tribal agency responsible for pesticide regulation.

#### 4.0 AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements in this labeling about Personal Protective Equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

#### 5.0 GENERAL INFORMATION FOR USE

**IMPORTANT:** Before application, read all use directions thoroughly. Use ReTain® only as directed. For specific information on the best use practices for your location, [consult with your Valent Agricultural Specialist \(1-800-6-Valent\)](#).

ReTain contains 15% (wt/wt) [S]-*trans*-2-amino-4-(2-aminoethoxy)-3-butenoic acid hydrochloride, commonly known as aminoethoxyvinylglycine, aviglycine HCl, or AVG HCl. AVG HCl is a potent plant growth regulator produced by fermentation. AVG HCl inhibits the production of ethylene in plant tissues. Ethylene affects plant processes such as flower development, fruit set, fruit maturation, fruit ripening, and fruit abscission.

#### 6.0 GENERAL APPLICATION INSTRUCTIONS

##### IMPORTANT

- Do not apply this product through any type of irrigation system.
- ReTain is packaged in a water-soluble bag sealed inside a metallized polyester pouch. Each water-soluble bag contains 11.7 oz (333 grams) of ReTain. Once the outer pouch is opened, the entire water-soluble bag must be used.
- Because the proper timing of ReTain application is particularly important to product performance, serious consideration is to be given, on a block-by-block basis, to any site conditions (e.g., elevation, sun exposure, soil texture), growing condition(s) (e.g., prevailing or anticipated weather patterns such as high temperature, chilling, drought or flood conditions, nutrient levels) or production practices (e.g., crop load, pruning severity) that could impact flowering, fruit set, or fruit maturation during the course of the season.

- On tree crops and cucumbers, apply ReTain in a sufficient amount of water to ensure that flowers, fruits, and foliage receive thorough spray coverage using calibrated spray equipment. Adjust water volumes based on plant size and spacing. However, excessive spray application volumes resulting in spray runoff will reduce product efficacy.
- Prepare spray solution by mixing the required amount of product with water in a clean, empty spray tank. Dispose of any unused spray material at the end of each day.
- Avoid applications during the heat of the day. For best results, apply ReTain under slow drying conditions, e.g., early in the morning or at night, in order to maximize absorption.
- Maintain solution pH between 6 and 8.
- For apple, pear, stone fruit, walnut, pecan and cucumber, do not apply ReTain to plants or fruit under stress (e.g., heat, water, disease, insect). Injured or stressed plants or fruits may not respond to ReTain.
- Do not use overhead cooling or irrigation equipment for at least 8 hours following ReTain application.
- Do not apply ReTain if rain is expected within 8 hours of application.

#### 7.0 COMPATIBILITY WITH OTHER AGRICULTURAL PRODUCTS

ReTain can be tank-mixed with DiPel®, Biobit®, and Xen-Tari® biological insecticides. Compatibility and performance data for ReTain with other agricultural products are not available.

Do not tank mix ReTain with other products unless compatibility has been verified. If considering tank mixing ReTain with other products use the following **compatibility jar test** before mixing a whole tank:

Add water from the same water source to a clear glass or plastic jar. Add the pesticides in correct proportions. Mix thoroughly and let stand for a minimum 15 minutes. Separation, gelling, or generation of heat are all signs of incompatibility.

Even if a mix passes the jar test for compatibility, it is imperative to test it on a designated area to evaluate for phytotoxicity or ineffectiveness.

Always read and follow all label directions and precautions of each product. When using combinations of products the most restrictive of label limitations and precautions must be followed. Do not mix with any pesticide that has a prohibition against tank mixing. For further information consult your Valent Agricultural Specialist (1-800-6 Valent).

8.0 APPLE AND PEAR – FOR HARVEST MANAGEMENT AND IMPROVEMENT OF FRUIT QUALITY

CROP	OBJECTIVE / BENEFIT	APPLICATION TIMING / USE INSTRUCTIONS
<p><b>Apple</b></p>	<p><b>Single Application:</b> Depending on cultivar, orchard conditions, application timing, and grower objectives, one or more of the following benefits will be associated with ReTain:</p> <ul style="list-style-type: none"> <li>• Delayed fruit maturity</li> <li>• Improved harvest management</li> <li>• Reduced preharvest fruit drop</li> <li>• Additional time for increase in fruit size</li> <li>• Additional time for color development</li> <li>• Maintenance of fruit firmness</li> <li>• Improved fruit quality (e.g., reduced incidence and/or severity of watercore and stem bowl cracking)</li> <li>• Enhanced storage potential</li> </ul>	<p><b>Single Pick Harvest:</b> Apply one to two pouches of ReTain per acre 21 to 28 days prior to the anticipated beginning of the normal harvest period of untreated fruit. ReTain applied 21 to 28 days before harvest will delay the harvest period up to 7 to 10 days. Applications made either too early or too late will significantly reduce the efficacy of the product.</p> <p><b>Multiple Pick Harvest:</b> To improve the fruit quality and storage potential of later picked apples (2nd, 3rd, 4th picks) apply one pouch of ReTain per acre, 7 to 14 days prior to the anticipated beginning of the normal harvest period of untreated fruit. ReTain applied 7 to 14 days before harvest typically will not delay the start of the harvest (1st pick), but will help control the maturation rate of the later picks.</p>
	<p><b>Double Application:</b> Depending on cultivar, orchard conditions, application timing, and grower objectives, one or more of the following benefits will be associated with a single application of ReTain:</p> <ul style="list-style-type: none"> <li>• Delayed fruit maturity</li> <li>• Improved harvest management</li> <li>• Reduced preharvest fruit drop</li> <li>• Additional time for increase in fruit size</li> <li>• Additional time for color development</li> <li>• Maintenance of fruit firmness</li> <li>• Improved fruit quality (e.g., reduced incidence and/or severity of watercore and stem bowl cracking)</li> <li>• Enhanced storage potential</li> </ul> <p>In addition to the effects of a single application of ReTain, one or more of the following benefits will be associated with a second application of ReTain:</p> <ul style="list-style-type: none"> <li>• Extended delay in fruit maturity</li> <li>• Extended harvest management period</li> <li>• Extended pre-harvest drop control</li> <li>• Extended maintenance of fruit quality at harvest</li> <li>• Greater enhancement of cold storage potential</li> </ul>	<p><b>For the First Application:</b> Apply one pouch of ReTain per acre 28 days prior to the anticipated beginning of the normal harvest period of untreated fruit.</p> <p><b>For the Second Application:</b> Apply one pouch of ReTain per acre 14 to 21 days after the first application but not less than 7 days before harvest.</p>
<p><b>Pear</b></p>	<p><b>Single Application:</b> ReTain will offer one or more of the same benefits as listed for apples, depending on the cultivar, growing conditions, and grower objectives.</p>	<p>Apply one pouch of ReTain per acre 7 to 16 days prior to the anticipated beginning of the normal harvest period of untreated fruit.</p>
	<p><b>Double Application:</b> ReTain will offer one or more of the same benefits as listed for apples, depending on the cultivar, growing conditions, and grower objectives.</p>	<p><b>For the First Application:</b> Apply one pouch of ReTain per acre 21 to 28 days prior to the anticipated beginning of the normal harvest period of untreated fruit.</p> <p><b>For the Second Application:</b> Apply one pouch of ReTain per acre 14 to 21 days after the first application but not less than 7 days prior to harvest.</p>
<p><b>Note:</b> Harvest management and fruit quality benefits from ReTain will be greatest when applied at one to two pouches per acre. Under difficult coloring conditions, color development of certain bi-color apple cultivars such as Gala and Honeycrisp™ will be delayed when ReTain is applied at one to two pouches per acre. For specific information on the best use practices for your location, consult with your Valent Agricultural Specialist (1-800-6-Valent).</p>		
<p><b>Water Volume:</b> ReTain efficacy requires that fruit and foliage receive thorough spray coverage. To ensure thorough coverage adjust water volumes based on tree size and spacing and use calibrated spray equipment (i.e., orchard air blast sprayer). Excessive spray application volumes that result in spray runoff will reduce product performance. In most cases, 100 gallons per acre has been shown to be effective.</p>		

(continued)

**APPLE AND PEAR – FOR HARVEST MANAGEMENT AND IMPROVEMENT OF FRUIT QUALITY** (continued)

<p><b>Adjuvants:</b> For optimal response, use ReTain with a 100% organosilicone surfactant. Use a final surfactant concentration of 0.05 to 0.1% (v/v) in the spray tank. To prevent possible spotting, use the 0.05% (v/v) concentration when high temperature (in excess of 86°F) weather conditions prevail or are anticipated. Do not use a surfactant concentration greater than 0.1% (v/v). To reduce foaming, add the adjuvant last and minimize agitation.</p>
<p><b>Harvest:</b> The normal harvest period for a particular orchard block refers to that time when fruits not treated with ReTain would be harvested for long term storage. To help determine the beginning of the normal harvest period, refer to historical trends for harvest dates and the “days from full bloom to harvest” interval for each cultivar in your area.</p>
<p><b>Additional Notes:</b></p> <ul style="list-style-type: none"> <li>• Apply ReTain before laying reflective film or foil mulch under ReTain treated apple trees. ReTain may be applied either before or after laying fabric groundcovers under ReTain treated apple trees.</li> <li>• ReTain has a 7 day preharvest interval (PHI) for Apple and Pear.</li> <li>• Do not tank-mix ReTain with sunburn protection product. If sunburn protection products are used, apply at least 3 days before or after ReTain application.</li> </ul>

**9.0 STONE FRUIT (EXCEPT CHERRY) – FOR HARVEST MANAGEMENT AND IMPROVEMENT OF FRUIT QUALITY**

CROP	OBJECTIVE / BENEFIT	APPLICATION TIMING / USE INSTRUCTIONS
<p><b>Apricot</b>  <b>Nectarine</b>  <b>Peach</b>  <b>Plum</b>  <b>Prune</b>  <b>(Interspecific variations of these crops, such as Plum-cot, pluot®, and Aprium®)</b></p>	<p>Depending on cultivar, orchard conditions, and grower objectives, one or more of the following benefits will be associated with ReTain:</p> <ul style="list-style-type: none"> <li>• Improved harvest management</li> <li>• Additional time for increase in fruit size</li> <li>• Maintenance of fruit firmness</li> <li>• Reduced preharvest fruit drop</li> <li>• Improved fruit quality</li> <li>• Enhanced storage potential</li> </ul>	<p>Apply one pouch of ReTain per acre one to two weeks prior to the anticipated beginning of the normal harvest period of untreated fruit.</p>
<p><b>Water Volume:</b> ReTain efficacy requires that fruit and foliage receive thorough spray coverage. To ensure thorough coverage adjust water volumes based on tree size and spacing and use calibrated spray equipment (i.e., orchard air blast sprayer). Excessive spray application volumes that result in spray runoff will reduce product performance. In most cases, 100 gallons per acre has been shown to be effective.</p>		
<p><b>Use of Adjuvants:</b> For optimal response, use ReTain with a 100% organosilicone surfactant. Use a final surfactant concentration of 0.05 to 0.1% (v/v) in the spray tank. To reduce foaming, add the adjuvant last and minimize agitation.</p>		
<p><b>Harvest:</b> The normal harvest period for a particular orchard block refers to that time when fruits not treated with ReTain would be harvested. To help determine the beginning of the normal harvest period, refer to historical trends for harvest dates and the “days from full bloom to harvest” interval for each cultivar in your area, and closely monitor the fruit maturity development for the current season.</p>		
<p><b>Additional Notes: Pre Harvest Interval (PHI):</b> ReTain has a 7 day (PHI) for labeled Stone Fruit.</p>		

**10.0 NUT, POME AND STONE FRUIT CROPS – FOR INCREASING FRUIT AND NUT SET**

CROP	OBJECTIVE / BENEFIT	APPLICATION TIMING / USE INSTRUCTIONS
<b>Apple</b>	To increase fruit set.	Apply one pouch of ReTain per acre, as a single application from pink stage to full bloom. Applications made prior to pink stage or after full bloom will significantly reduce efficacy of the treatment. Do not apply after petal fall.
<b>Cherry</b>	To increase fruit set.	Apply one to two pouches of ReTain per acre during bloom. Retain may be applied as a single application of up to two pouches, or as sequential applications of one pouch per application. Applications between popcorn stage (balloon stage) to first bloom are more effective than earlier or later applications. Do not apply after petal fall.
<b>European Pear cultivars</b>	To increase fruit set.	Apply one pouch of ReTain per acre, as a single application from white bud stage to full bloom. Applications made prior to white bud stage or after full bloom will significantly reduce efficacy of the treatment. Do not apply after petal fall.
	To reduce June drop.	Apply one pouch of ReTain at 10 mm fruit size.

(continued)

**NUT, POME AND STONE FRUIT CROPS – FOR INCREASING FRUIT AND NUT SET (continued)**

CROP	OBJECTIVE / BENEFIT	APPLICATION TIMING / USE INSTRUCTIONS
<b>Pecan</b>	To increase nut set in cultivars that experience low set.	Apply one to two pouches of ReTain per acre during bloom. Retain may be applied as a single application of up to two pouches, or as sequential applications of one pouch per application. Applications made either too early (pre-bloom) or too late (after full bloom) will reduce the efficacy of the treatment. Use higher rates on cultivars/orchards that have experienced lower nut set.
<b>Walnut</b>	Increased nut set in cultivars affected by pistillate flower abortion (PFA).	Apply one to two pouches of ReTain per acre, when most pistillate (female) flowers are receptive to pollination. ReTain may be applied as a single application of up to two pouches, or as sequential applications of one pouch per application. One application of ReTain at 5-30% pistillate flower bloom has been shown to be effective for a short pistillate flower bloom period. Sequential applications at 5-30% and 45-65% bloom are effective and cover a broader window of pistillate flower bloom. Applications made either too early (pre-bloom) or too late (full bloom or later) will significantly reduce efficacy of the treatment. Use higher rates on cultivars/orchards that have experienced high incidences of poor nut set due to PFA.
<p><b>IMPORTANT:</b></p> <ul style="list-style-type: none"> <li>• For apple and pear, do not apply more than one pouch for increasing fruit set (apples and pears) and June drop (pears). Do not use more than two pouches for harvest management and improvement of fruit quality. Do not apply more than two pouches of ReTain per acre per year.</li> <li>• For sweet cherry, do not apply more than one to two pouches for increasing fruit set. Do not apply ReTain to cherry trees when fruit is present.</li> <li>• For pecan and walnut, do not apply more than two pouches of ReTain per acre per year.</li> </ul>		
<p><b>Water Volume:</b> ReTain efficacy requires that flower buds and flowers receive thorough spray coverage. To ensure thorough coverage use calibrated spray equipment (i.e., air blast sprayer) and adjust water volumes based on tree size and spacing. Use sufficient water volume to achieve thorough coverage without excessive runoff. In most cases, a spray volume of 100 to 200 gallons per acre has been shown to be effective.</p>		
<p><b>Use of Adjuvants:</b> Use of adjuvants with ReTain for bloom applications has not been fully researched. Some adjuvants will cause a bloom thinning effect and will reduce the benefits of ReTain.</p>		

**11.0 CUCUMBER – FOR SEED PRODUCTION ONLY**

CROP	OBJECTIVE / BENEFIT	APPLICATION TIMING / USE INSTRUCTIONS																
<b>Cucumber</b> Gynoecious inbred breeding lines.	Promotion of staminate (male) flowers in gynoecious (female flowering) cucumber-breeding lines for the purpose of producing inbred and hybrid cucumber seed.	Apply ReTain directly to gynoecious plants to be converted to staminate flowers. Make one to six applications of 250 ppm to 1000 ppm AVG HCL beginning at early first leaf stage through tenth leaf stage. Use the appropriate skip row technique to achieve desired sex ratio of plants.																
<p><b>Water Volume:</b> Apply ReTain in a sufficient amount of water to ensure good thorough coverage. In most cases, 26 to 52 gallons per treated acre will be sufficient. The application rate and frequency of application will depend on the sensitivity of the individual inbred line to ReTain. Leaf yellowing can occur at high rates. If unacceptable leaf yellowing is noted, use lower concentrations in subsequent applications.</p>																		
<p><b>Dilution Table:</b> Parts Per Million (ppm) AVG HCl with varying Gallons (gal) of spray solution and pouches of ReTain for cucumber use.</p>																		
<table border="1"> <thead> <tr> <th data-bbox="363 1581 602 1625">Gal Spray Solution</th> <th data-bbox="602 1581 841 1625">Pouches of ReTain</th> <th data-bbox="841 1581 1079 1625">Grams AVG HCl</th> <th data-bbox="1079 1581 1317 1625">PPM AVG HCl</th> </tr> </thead> <tbody> <tr> <td data-bbox="363 1625 602 1669">13</td> <td data-bbox="602 1625 841 1669">1</td> <td data-bbox="841 1625 1079 1669">50</td> <td data-bbox="1079 1625 1317 1669">1000</td> </tr> <tr> <td data-bbox="363 1669 602 1713">26</td> <td data-bbox="602 1669 841 1713">1</td> <td data-bbox="841 1669 1079 1713">50</td> <td data-bbox="1079 1669 1317 1713">500</td> </tr> <tr> <td data-bbox="363 1713 602 1753">52</td> <td data-bbox="602 1713 841 1753">1</td> <td data-bbox="841 1713 1079 1753">50</td> <td data-bbox="1079 1713 1317 1753">250</td> </tr> </tbody> </table>			Gal Spray Solution	Pouches of ReTain	Grams AVG HCl	PPM AVG HCl	13	1	50	1000	26	1	50	500	52	1	50	250
Gal Spray Solution	Pouches of ReTain	Grams AVG HCl	PPM AVG HCl															
13	1	50	1000															
26	1	50	500															
52	1	50	250															
<p><b>Use of Adjuvants:</b> Addition of an adjuvant to ReTain increases the likelihood of leaf yellowing in young cucumber plants.</p>																		
<p><b>Additional Note:</b> Do not reapply ReTain within 24-hours of application. Use of ReTain on cucumber is for gynoecious breeding line and hybrid seed production only. Do not apply directly to, or allow spray to drift to cucumber crops grown for fresh or processed markets. Fruit from plants treated with ReTain is not to be used for human or animal consumption.</p>																		

12.0

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store product in a cool and dry place. Avoid extended storage conditions at temperatures above 25°C (77°F).

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty container into applicable equipment. If recycling is not available dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

13.0 **NOTICE TO USER**

**Warranty and Disclaimer Statement:**

To the extent consistent with applicable law, seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning the use of this product other than as indicated on the label. To the extent consistent with applicable law, user assumes all risks of use, storage or handling not in strict accordance with accompanying directions.

Biobit®, DiPel®, and XenTari® are registered trademarks of Valent BioSciences LLC.

*Products That Work, From People Who Care* is a registered trademark of Valent U.S.A. LLC.

Aprium® and Pluot® are registered trademarks of Zaiger Genetics.

Honeycrisp™ is a registered trademark of the University of Minnesota.

Registered by:

**Valent BioSciences LLC**

870 Technology Way  
Libertyville, IL 60048 U.S.A.

List No. 12017-04  
Form 04-9675/R17



For state registration and/or supplemental labels, please call or visit us online.  
*Products That Work, From People Who Care*® | [valent.com](http://valent.com) | 800-6-VALENT (682-5368)  
**Always read and follow label instructions.**

©2017 Valent BioSciences LLC. All rights reserved. Printed in the U.S.A.