

Allium Scouting Form						
Farm/Grower:	Scout:					
Field Name:	Date:					
Cron/Cultivar:	Block:					

Crop Stage: < 3 leaf Bulb enlargement Leaf senesence Harvest

Procedure: Select plants at random by using the same # of paces between plants to cover a field or block, making 10 total stops. Inspect 3 plants in a row at each stop.

Thrips: Begin scouting in early June. Pry apart the leaves gently counting the number of thrips and leaves per plant. Inspect plants up to the tips of the leaves.

Leek Moth: Begin scouting in May. Record presence or absence. Larvae are yellowish-green with a pale brown head capsule and 8 small grey spots on each abdominal segment. At maturity, larvae reach 13-14 mm in length.

Other Insect: Pick your own ____ = ___

Insect Damage: Ch: Chewing can be symptoms of Leek moth larvae feeding **Sc:** scratching marks from thrips feeding **Tw:** Leaf twisting from Thrips feeding. **Be:** Plants bending from thrips feeding.

Disease ID: PB: Purple blotch Bac: Sour Skin or Bacterial soft rot Bot: Botrytis DM: Downy Mildew. Indicate if sample taken for ID.

Plant	# of leaves per plant	# of Thrips per plant	Leek Moth	Other insect	Insect Damage (Ch, Sc, Tw, Be)	Disease ID (PB, Bac, Bot, DM)	% Leaves Affected by Disease	Other Notes/Weeds
1								
2								
3								
4								
5								
6								
7								
8								
9								
10								
11								
12								
13								
14								
15								
16								
17								
18								
19								
20								
21								
22								
23								
24								
25								
26								
27								
28								
29								
30								
Total								
AVG.								

Scouting Map